



## King's Research Portal

DOI:

[10.1016/j.nicl.2016.06.020](https://doi.org/10.1016/j.nicl.2016.06.020)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Wolfers, T., Van Rooij, D., Oosterlaan, J., Heslenfeld, D., Hartman, C. A., Hoekstra, P. J., Beckmann, C. F., Franke, B., Buitelaar, J. K., & Marquand, A. F. (2016). Quantifying patterns of brain activity: Distinguishing unaffected siblings from participants with ADHD and healthy individuals. *NeuroImage: Clinical*.  
<https://doi.org/10.1016/j.nicl.2016.06.020>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## Accepted Manuscript

Quantifying patterns of brain activity: Distinguishing unaffected siblings from participants with ADHD and healthy individuals; QUERY id="Q3" name="IGALVEZ" Some address was captured as footnote. Please if appropriate and correct if necessary. QUERY



Thomas Wolfers, Daan van Rooij, Jaap Oosterlaan, Dirk Heslenfeld, Catharina A. Hartman, Pieter J. Hoekstra, Christian F. Beckmann, Barbara Franke, Jan K. Buitelaar, Andre F. Marquand

PII: S2213-1582(16)30118-8  
DOI: doi: [10.1016/j.nicl.2016.06.020](https://doi.org/10.1016/j.nicl.2016.06.020)  
Reference: YNICKL 763

To appear in: *NeuroImage: Clinical*

Received date: 20 May 2016  
Revised date: 24 June 2016  
Accepted date: 28 June 2016

Please cite this article as: Wolfers, Thomas, van Rooij, Daan, Oosterlaan, Jaap, Heslenfeld, Dirk, Hartman, Catharina A., Hoekstra, Pieter J., Beckmann, Christian F., Franke, Barbara, Buitelaar, Jan K., Marquand, Andre F., Quantifying patterns of brain activity: Distinguishing unaffected siblings from participants with ADHD and healthy individuals; QUERY id="Q3" name="IGALVEZ" Some address was captured as footnote. Please if appropriate and correct if necessary. QUERY, *NeuroImage: Clinical* (2016), doi: [10.1016/j.nicl.2016.06.020](https://doi.org/10.1016/j.nicl.2016.06.020)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Quantifying patterns of brain activity: distinguishing unaffected siblings from participants with ADHD and healthy individuals

Running title: Quantifying patterns of brain activity

Thomas Wolfers<sup>1,2,\*</sup>, Daan van Rooij<sup>2</sup>, Jaap Oosterlaan<sup>3</sup>, Dirk Heslenfeld<sup>3</sup>, Catharina A. Hartman<sup>4</sup>, Pieter J. Hoekstra<sup>4</sup>, Christian F. Beckmann<sup>2,5,6</sup>, Barbara Franke<sup>1,7</sup>, Jan K. Buitelaar<sup>5,8</sup>, Andre F. Marquand<sup>2,9</sup>

\*corresponding author

<sup>1</sup> Department of Human Genetics, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, Nijmegen, The Netherlands

<sup>2</sup> Donders Centre for Cognitive Neuroimaging, Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

<sup>3</sup> Clinical Neuropsychology section, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

<sup>4</sup> University of Groningen, University Medical Center Groningen, Groningen, Department of Psychiatry, The Netherlands

<sup>5</sup> Department of Cognitive Neuroscience, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Centre, Nijmegen, The Netherlands

<sup>6</sup> Centre for Functional MRI of the Brain (FMRIB), University of Oxford, Oxford, United Kingdom

<sup>7</sup> Department of Psychiatry, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, Nijmegen, The Netherlands

<sup>8</sup> Karakter Child and Adolescent Psychiatry University Centre, Radboud University Medical Centre, Nijmegen, The Netherlands

<sup>9</sup> Department of Neuroimaging, Institute of Psychiatry, King's College London, United Kingdom

Number of words abstract: 224 words

Number of words article (excluding abstract, acknowledgements, financial disclosures sections, legends, and references): 3559 words

Number of tables: 1

Number of figures: 2

Number of supplementary material: 4

*Thomas Wolfers (corresponding author)*

Donders Centre for Cognitive Neuroimaging  
Radboud University  
PO Box 9101  
6500 HB Nijmegen  
The Netherlands  
Phone: +31-24-3610098  
E-mail: t.wolfers@donders.ru.nl

*Dr. Daan van Rooij*

Donders Centre for Cognitive Neuroimaging  
Radboud University  
PO Box 9101  
6500 HB Nijmegen  
The Netherlands  
E-mail: D.vanRooij@donders.ru.nl

*Prof. Dr. Jaap Osterlaan*

Clinical neuropsychology  
Vrije Universiteit Amsterdam  
De Boelelaan 1105,  
1081 HV Amsterdam,  
The Netherlands  
E-mail: j.oosterlaan@vu.nl

*Dr. Dirk Heslenfeld*

Cognitive Neuropsychology  
Vrije Universiteit Amsterdam  
De Boelelaan 1105,  
1081 HV Amsterdam,  
The Netherlands  
E-mail: d.j.heslenfeld@vu.nl

*Dr. Catharina A. Hartman*

Department of Psychiatry  
University Medical Center Groningen  
University of Groningen  
Hanzeplein 1,  
9713 GZ Groningen  
The Netherlands  
E-mail: c.hartman@accare.nl

*Prof. Dr. Pieter J. Hoekstra*

Department of Psychiatry  
University Medical Center Groningen  
University of Groningen  
Hanzeplein 1,  
9713 GZ Groningen  
The Netherlands  
E-Mail: p.hoekstra@accare.nl

*Prof. Dr. Christian F. Beckmann*

Donders Centre for Cognitive Neuroimaging  
Radboud University  
Postbus 9101  
6500 HB Nijmegen  
The Netherlands  
E-Mail: c.beckmann@donders.ru.nl

*Prof. Dr. Barbara Franke*  
Department of Human Genetics  
Radboud University Medical Center  
PO Box 9101  
6500 HB Nijmegen  
The Netherlands  
E-Mail: b.franke@donders.ru.nl

*Prof. Dr. Jan K. Buitelaar*  
Karakter Child and Adolescent Psychiatry University Centre  
Radboud University Medical Centre  
PO Box 9101  
6500 HB Nijmegen  
The Netherlands  
E-Mail: jan.buitelaar@radboudumc.nl

*Dr. Andre F. Marquand*  
Donders Centre for Cognitive Neuroimaging  
Radboud University  
PO Box 9101  
6500 HB Nijmegen  
The Netherlands  
E-Mail: a.marquand@donders.ru.nl

**Abstract**

Attention-deficit/ hyperactivity disorder (ADHD) is one of the most prevalent and heritable psychiatric disorders. While previous studies have focussed on mapping focal or connectivity differences at the group level, the present study employed pattern recognition to quantify group separation between unaffected siblings, participants with ADHD, and healthy controls on the basis of spatially distributed brain activations. This was achieved using an fMRI-adapted version of the Stop-Signal Task in a sample of 103 unaffected siblings, 184 participants with ADHD, and 128 healthy controls. We used activation maps derived from three task regressors as features in our analyses employing a Gaussian process classifier. We showed that unaffected siblings could be distinguished from participants with ADHD (area under the receiver operating characteristic curve (AUC) = .65,  $p=.002$ , 95% Modified Wald CI: .59 - .71 AUC) and healthy controls (AUC = .59,  $p=.030$ , 95% Modified Wald CI: .52 - .66 AUC), although the latter did not survive correction for multiple comparisons. Further, participants with ADHD could be distinguished from healthy controls (AUC = .64,  $p=.001$ , 95% Modified Wald CI: .58 - .70 AUC). Altogether the present results characterise a pattern of frontolateral, superior temporal and inferior parietal expansion that is associated with risk for ADHD. Unaffected siblings show differences primarily in frontolateral regions. This provides evidence for a neural profile shared between participants with ADHD and their healthy siblings.

Keywords: Pattern recognition, Gaussian process classification, fMRI, Response inhibition, Attention - deficit/ hyperactivity disorder, Unaffected Siblings,

## Introduction

Attention-deficit/ hyperactivity disorder (ADHD) is one of the most prevalent (Polanczyk et al., 2007; Simon et al., 2009) and heritable psychiatric disorders (Faraone and Biederman, 2005). Heritability estimates are around 75%, and relatives of participants with ADHD have about 25% risk to have ADHD themselves which is about four times higher than the population rate (Biederman et al., 1990). While unaffected siblings share some of the biological risk for disease with their affected siblings, they do not express this risk symptomatically. Unaffected siblings of participants with ADHD have shown patterns of cognitive and neural functioning intermediate to those observed in affected siblings and healthy controls (Durstion et al., 2006; Greven et al., 2015; Slaats-Willemse et al., 2003). Specifically, unaffected siblings of participants with ADHD have shown impairments in response inhibition, as evidenced in impaired performance on cognitive tasks, as well as brain abnormalities in structures subserving response inhibition. Therefore, this group of individuals represents an interesting study population in exploring disorder mechanisms and a complex group for disorder classification as preformed in this study. Response inhibition deficits belong to the most prevalent deficits observed in children and adolescents with ADHD (Barkley, 1999; Castellanos et al., 2006; Hart et al., 2014a; Slaats-Willemse et al., 2003; van Rooij et al., 2015b). To probe deficits in response inhibition in the current study, we chose the Stop-Signal Task functional magnetic resonance imaging (fMRI) paradigm. This experimental paradigm has previously shown reductions in task related brain (de)activations in participants with ADHD compared to healthy controls (Hart et al., 2014a; Janssen et al., 2015; Lipszyc and Schachar, 2010; van Rooij et al., 2015b).

In contrast to studies that examined focal or connectivity differences between unaffected siblings and participants with ADHD (van Rooij et al., 2015a, 2015b), we quantified group separation on the basis of spatially distributed patterns of activity across the brain, which provides a unified measure of group separation that is more representative of the overall pattern of brain activity than any individual region. Pattern recognition is ideal for this purpose and aims to extract regularities in data, which can be used to predict group membership (Hastie et al., 2009). Early pattern recognition studies aimed to show that participants with ADHD could be distinguished from healthy controls based on different MRI modalities (Hart et al., 2014a, 2014b; Igual et al., 2012; Lim et al., 2013; Wang et al., 2011). These studies were usually small in size and the literature tends to show reduced classification performance with increased sample size (Wolfers et al., 2015). Larger studies capture more of the inherent heterogeneity of ADHD, in terms of its symptomatology and pathophysiology. Therefore, those studies are more indicative for the predictability of ADHD in clinical settings, as a heterogeneous group of patients approach clinics to seek treatment.

In studies on unaffected siblings of schizophrenia and autism spectrum disorders, researchers used neural patterns to distinguish siblings from their respective patient group and healthy controls (Fan et al., 2008; Segovia et al., 2014; Yu et al., 2013). However, until now no pattern recognition study has investigated unaffected siblings of participants with ADHD. In the present study we sought to: (i) precisely quantify the group separation between unaffected siblings, participants with ADHD and healthy controls in a large sample that accurately reflects the range of variation in the disease phenotype and (ii) map the nature of these differences to identify response inhibition related activation patterns, that underlay the shared genetic load between unaffected siblings and participants with ADHD. The present study is the largest study employing



pattern recognition to investigate unaffected siblings of participants with ADHD, using a hallmark deficit of ADHD as biomarker, response inhibition (Barkley, 1999; Castellanos et al., 2006; Hart et al., 2014a; Slaats-Willemse et al., 2003; van Rooij et al., 2015b).

## Methods

### *Participants*

We used data from the NeuroIMAGE project, a large longitudinal clinical cohort consisting of individuals tested at two different sites in The Netherlands, the Vrije Universiteit in Amsterdam and the Donders Centre for Cognitive Neuroimaging in Nijmegen. We selected all individuals who performed the Stop-Signal Task. ADHD diagnosis was based on K-SADS (Birmaher et al., 2010) structured psychiatric interviews and Conners' questionnaires (Conners et al., 1998). The total sample consisted of 184 participants with ADHD, 103 unaffected siblings, and 128 healthy controls (Table 1). This sample is similar to the sample detailed in our previous publication (van Rooij et al., 2015b), with the exception that the current study excluded subjects if there was an inconsistent diagnosis based on either K-SADS or Conners' questionnaire. Ethics approval for this study was obtained from relevant ethics review boards, and informed consent/assent was signed by parents and their children. A comprehensive overview of recruitment, diagnostics, ethical approval, testing procedures, and quality control are provided in a separate methods publication (von Rhein et al., 2014).

*[Table 1: Demographic and clinical characteristics of complete sample]*

### Stop-Signal Task design

Response inhibition was measured using an fMRI-adapted version of the Stop-Signal Task (van Meel et al., 2007; van Rooij et al., 2015b), consisting of four blocks of 60 trials each. Participants were instructed to respond as quickly and accurately as possible to a go-signal (two-choice reaction time task) with a left or right button press

on a button box, unless the go-signal was followed by a stop-signal (25% of trials), in which case participants were instructed to withhold their response. Participants who did not reach 70% accuracy on the go-trials were excluded prior to analyses (N=5). The task was adapted to the performance of the participant, by varying the delay between go and stop-signal (stop-signal delay), in order to achieve 50% successful inhibition on stop-trials for all participants. The stop-signal delay was decreased from an initial 250ms, by 50ms after successful inhibition, and increased by 50ms after failed inhibition. The main measure of response inhibition performance, the stop-signal reaction time (SSRT), was calculated by averaging the delay necessary for a participant to successfully inhibit his/her response in 50% of the stop-trials. Secondary outcome measures were the total number of omission and commission errors on go-trials (errors) and the intra-individual component of variation (ICV), calculated by dividing the reaction time variance by the mean reaction time (both calculated from reaction times on correct go trials).

#### *Acquisition of functional MRI*

Data were acquired at both sites on similar 1.5 Tesla Siemens MRI scanners (Siemens Sonata at VUmc; Siemens Avanto at Donders Centre for Cognitive Neuroimaging) using the same Siemens 8-channel head coil and the following protocol: The Stop-Signal Task was collected in four runs using a T2\*-weighted echo planar imaging sequence (TR=2340 ms, TE=40 ms, FOV=224x224 mm, 37 slices, voxel size=3.5x3.5x3.5 mm, 94 volumes per run). To assist accurate normalization, participants were also scanned using a high resolution MPRAGE T1-weighted sequence (TR=2730 ms, TE=2.95 ms, TI=1000 ms, flip angle = 7°, voxel size=1x1x1 mm, matrix size = 256 × 256, FOV=256 mm, 176 slices).

### *Processing of fMRI data*

Functional MRI data were processed using FSL (FMRIB's Software Library, [www.fmrib.ox.ac.uk/fsl](http://www.fmrib.ox.ac.uk/fsl); fMRI Expert Analysis Tool, version 6.0 (Jenkinson et al., 2012; Smith et al., 2004; Woolrich et al., 2009). Preprocessing included removal of the first four volumes of each run, within-run motion correction to the middle volume, slice-timing correction, and spatial smoothing with a 6 mm Gaussian kernel, before residual motion correction was applied using ICA-AROMA. ICA-AROMA is an advanced motion correction tool that has been shown to outperform other motion correction procedures (Pruim et al., 2015a, 2015b). The data from each run were registered to the participant's T1 anatomical image using linear, boundary-based registration implemented in FSL-FLIRT. For each participant a general linear model was fit, including successful stop, failed stop and successful go trials as regressors in addition to error trials, signal from cerebral spinal fluid and white matter, which were included as nuisance covariates. Task regressors were convolved with a double-gamma hemodynamic response function and data were high pass filtered with a cutoff of .01 Hz prior to estimation. The resulting single-subject regression coefficient images ('beta maps') were transformed to participant-level anatomical space (3 mm isotropic resolution) and combined across runs using a fixed effects model, using FSL-FEAT. This resulted in three participant-level activation maps, (1) successful stop, (2) failed stop, and (3) successful go, which were transformed to a neutral 'midspace', a procedure which neutralizes potential registration biases due to structural group and gender differences. The reader is referred to a prior publication for further details of the processing procedures (van Rooij et al., 2015b), where the only difference in the

present manuscript was the addition of the advanced motion correction using ICA-AROMA.

### *Quantifying and mapping group separation with Gaussian process classifiers*

Gaussian process classifiers (Rasmussen and Williams, 2006) were used to distinguish participants with ADHD from their unaffected siblings and healthy controls. Gaussian processes are best described as a distribution over functions, where inference proceeds by first computing the posterior distribution over functions according to the rules of probability. This is referred to as conditioning the prior distribution on the data. In the classification case, the posterior process is then passed through a sigmoid response function that maps the output to the unit interval, thereby providing a valid probability score for each prediction. These quantify predictive confidence and provide the primary advantage of Gaussian process classifiers over alternative approaches. Further details surrounding this approach have been published previously (Marquand et al., 2010). First, we estimated group separation on the basis of neuroimaging biomarkers. For this, we trained GPC models to make predictions based on the activation maps corresponding to the three task regressors, described in the fMRI processing section. The total number of features in these classifications was 224781. Second, we estimated group separation on the basis of behavioural data, which provides a reference for the classifier above. For this, we trained a GPC model on the basis of data from the behavioural task. Specifically, we used the number of errors during the task, the ICV as well as the SSRT as features (see above). Each classifier was embedded within a leave-one-participant-out cross-validation procedure, and the measure of generalizability was the area under the receiver-operating characteristic curve (AUC). This measure has the advantage that it is not sensitive to a particular choice of decision

threshold. Statistical significance was assessed by permutation testing for the AUC, taking into account the family structure within the sample. Specifically, instead of permuting the labels individually, we permuted the labels that belong to participants from the same family together. In that way, we ensured that the family structure was preserved, when the labels were shuffled.

Multiple-comparison correction for the AUCs was performed with the Bonferroni-Holm method (Holm, 2010) and 95% confidence intervals were reported and based on the modified Wald-method (Kottas et al., 2014). Note that these confidence intervals should be considered illustrative only. The primary measures we use to assess statistical significance are p-values derived from the permutation testing procedures described above, which fully account for the family structure in the data.

A common approach to visualising the importance of each brain region to the classification is to visualise the classifier weights directly (Mourao-Miranda et al 2005). However, the classifier weights are influenced by both the signal and noise in the data, which complicates interpretation. Therefore, forward maps (Haufe et al., 2014) were computed that provide a better indication of the differential activation pattern underlying the group separation. Most commonly, these maps are reported without applying a threshold, but it is clearly desirable to localise the most important differences. Therefore, we present a novel approach to thresholding forward maps based on fitting a mixture model. To achieve this, we fit a Gaussian-Gamma mixture to the image histograms that provide an explicit model for the null distribution plus positive and negative activations (Beckmann and Smith, 2004). For this, we used the implementation in the FSL-MELODIC software<sup>1</sup>. After fitting this model, these maps can then be thresholded in two ways: (i) by an alternative hypothesis testing (AHT)

---

<sup>1</sup> <http://fsl.fmrib.ox.ac.uk/fsl/fslwiki/MELODIC>

procedure where voxels are declared significant if they have a probability  $p_{\text{AHT}} > .5$  of belonging to one of the alternative distributions (Beckmann and Smith, 2004) or (ii) by controlling the false discovery rate (FDR) against the explicitly modelled null distribution (Efron, 2004; Efron et al., 2001). Here this was done at the nominal rate of  $p_{\text{FDR}} < .05$ . In our data, both approaches lead to similar conclusions (see supplementary figure 1). All figures were visualized in Caret (Van Essen et al., 2001).

Sensitivity analyses were performed to increase the confidence in the analyses described above. Since the total sample showed a slight class imbalance with respect to gender and scan-site and in order to reduce nuisance variance, the sample was perfectly matched for gender as well as scan-site and optimally on age. We used optimal matching algorithms implemented in the R-package MatchIt to simultaneously match age across all groups (Ho et al., 2011; for information on the matched sample see Supplementary Table 1).

The matched sample contained 74 participants per group. The analyses were performed in MATLAB using customized scripts from the PRoNTo toolbox (Schrouff et al., 2013).

## Results

### *Descriptive statistics: Stop-Signal Task*

Unaffected siblings showed shorter stop-signal reaction times and lower error rates than the participants with ADHD (Table 1, Wald  $\chi^2 = 7.941$ ,  $p < .005$  and Wald  $\chi^2 = 10.701$ ,  $p < .001$ , respectively), but did not differ from healthy controls in their reaction times and error rates (Table 1, Wald  $\chi^2 = .743$ ,  $p = .389$  and Wald  $\chi^2 = .5954$ ,  $p = .343$ , respectively). The intra-individual component of variation was lower in unaffected siblings as compared to the participants with ADHD (Table 1, Wald  $\chi^2 = 20.213$ ,  $p < .001$ ), and slightly higher than in healthy controls (Wald  $\chi^2 = 4.057$ ,  $p = .044$ ). These results are similar to those reported in an earlier study with an overlapping sample (van Rooij et al., 2015b). The addition of age, gender, IQ, medication status, or comorbid diagnoses to the model did not influence the reported group differences.

### *Quantifying and mapping group separation*

The accuracies for discriminating groups are summarised in Figure 1. Briefly, unaffected siblings could be distinguished from the participants with ADHD on the basis of successful stop activation maps (AUC = .65,  $p < .002$ , 95% Modified Wald CI: .59 - .71 AUC) and participants with ADHD could be distinguished from healthy controls on the basis of the same activation maps included as features to the classifications (AUC = .64,  $p < .001$ , 95% Modified Wald CI: .58 - .70 AUC). We also found nominally significant discrimination of unaffected siblings from healthy controls based on successful stop activation maps (AUC = .59,  $p < .030$ , 95% Modified Wald CI: .52 - .66 AUC) as well as nominally significant discrimination between participants with ADHD and healthy



controls based on failed stop activation maps ( $AUC = .60$ ,  $p < .019$ , 95% Modified Wald CI: .54-.66 AUC, Figure 1; for balanced accuracy, sensitivity and specificity measures in the complete and matched sample see Supplementary Table 2).

Figure 2 shows the forward maps for the successful stop activation maps for each group distinction without applying a threshold. In Supplementary Figure 1, the same maps are shown without a threshold in the first column, with a threshold of  $p_{AHT} > .5$  in the second and  $p_{FDR} < .05$  in the third. The classifier discriminating participants with ADHD from healthy controls showed a frontolateral, superior temporal and inferior parietal pattern with positive coefficients favouring ADHD. The pattern that separated unaffected siblings from participants with ADHD showed high coefficients in frontolateral and inferior parietal areas favouring ADHD, but was for the remainder wide-spread in comparison. The nominally significant unaffected sibling versus healthy controls distinction, showed a pattern with high coefficients primarily in inferior frontolateral areas favouring unaffected siblings. In Supplementary Figure 2 we show the fit of the mixture models to the three forward maps. The sensitivity analyses for which we perfectly matched the sample on gender and scan site and optimally on age showed a similar pattern of results as those described above, with exception of the successful-stop difference between ADHD and their unaffected siblings, all predictions improved in the matched sample (Figure1).

For the classifier trained to separate groups on the basis of the behavioural data, we showed that unaffected siblings could be distinguished from participants with ADHD ( $AUC = .66$ ,  $p < .001$ ) but not from healthy controls ( $AUC = .51$ ,  $p > .05$ ), based on behavioural scores described earlier. Participants with ADHD could be distinguished from healthy controls ( $AUC = .71$ ,  $p < .001$ ).

*[Figure 1; Figure 2]*

ACCEPTED MANUSCRIPT

## Discussion

In this study we showed that: (i) unaffected siblings of participants with ADHD could be distinguished from healthy controls and from participants with ADHD. Further, (ii) participants with ADHD could be reliably distinguished from controls. (iii) The predictions on behavioural data were approximately equally accurate, except for the distinction of unaffected siblings from healthy controls, which was not possible with behavioural data. The pattern of difference between participants with ADHD and healthy controls was characterised by positive bilateral frontolateral, superior temporal and inferior parietal coefficients favouring ADHD and frontolateral coefficients favouring unaffected siblings in comparison with healthy controls. This provides evidence for a neural profile shared between participants with ADHD and their healthy siblings.

The pattern of difference reported here, partially overlaps with regions in frontal and parietal areas reported in earlier studies of our NeuroImage sample (van Rooij et al., 2015a, 2015b). Looking at the thresholded forward maps (Supplementary Figure 1), we see frontolateral areas with positive coefficients favouring participants with ADHD as well as unaffected siblings when contrasted with healthy controls. In comparison to our previous studies that examined focal or connectivity differences, we extend these findings by precisely quantifying group separations based on task activation maps and show that the pattern of difference that distinguishes all groups is characterized by a widespread profile.

The diagnostic accuracy we report is moderate in relation to earlier studies aiming to separate participants with ADHD from controls using small samples (Wolfers et al., 2015) but is comparable to studies that have employed large samples that capture

more of the heterogeneity in the ADHD phenotype (Sabuncu and Konukoglu, 2014). The pattern recognition approach we employed allowed us to quantify the degree of separation between groups and therefore also the degree to which shared familial risk factors present in patients and unaffected siblings are expressed in patterns of brain activity. We could distinguish unaffected siblings from participants with ADHD showed a similar accuracy for distinguishing unaffected siblings from healthy controls based on fMRI data. This is in line with earlier studies that identified patterns of shared risk between siblings of participants with autism and schizophrenia (Fan et al., 2008; Segovia et al., 2014; Yu et al., 2013). Interestingly, a distinction based on behavioural data was not possible between unaffected siblings and healthy controls, indicating that unaffected siblings are not behaviourally different from healthy controls in response inhibition. However, they show a different neural pattern, which may be linked to compensatory brain processes in these unaffected individuals compared to their affected siblings.

As mentioned in the introduction, ADHD has mostly been classified in considerably smaller studies (Hart et al., 2014a, 2014b; Igual et al., 2012; Johnston et al., 2014; Lim et al., 2013; Peng et al., 2013; The ADHD Consortium, 2012; Wang et al., 2013, 2011; Zhu et al., 2008). In a classical analytic setting, p-values derived from measures of central tendency (e.g. a t-test) have an explicit dependency on the sample size, so the significance necessarily increases with increasing sample size, even though the effect size may not. In contrast, the predictive accuracy is a measure of class overlap that is governed by the distributions of the different classes and is largely independent of sample size, if properly assessed (e.g. using cross-validation). Therefore, the estimate of class overlap becomes more precise with increased sample size. This is important because the present study is the largest task-based fMRI study employing pattern

recognition in ADHD and therefore may represent a benchmark for what is possible in terms of accuracies in representative cohorts of heterogeneous disorders. This heterogeneity may, for example, stem from sampling subjects at different ages and at different points on their developmental trajectory. Our results suggest that – like all psychiatric disorders – the heterogeneity of the ADHD phenotype presents a major challenge for identifying disease mechanisms and for finding biomarkers that predict diagnosis and disease course. For example, previous research shows that only a subset of participants with ADHD display behavioural alterations in response inhibition (Mostert et al., 2015a, 2015b). Different participants with ADHD may have different symptom profiles and different underlying biological causes (Faraone et al., 2015). Therefore, finding methods to parse heterogeneity is a major research initiative. Clustering methods are most commonly used for this purpose and aim to partition patients into subgroups (Fair et al., 2012; Mostert et al., 2015b; van Hulst et al., 2014), but alternative methods such as normative modelling (Marquand et al., 2016a, 2016b) may also be beneficial for understanding heterogeneity underlying psychiatric disorders.

In summary, the present results describe a pattern of frontolateral, superior temporal and inferior parietal expansion that is associated with risk for ADHD. Unaffected siblings show differences primarily in frontolateral regions. This provides evidence for a neural profile shared between participants with ADHD and their healthy siblings. In the future, pattern recognition techniques can be employed to break down heterogeneity in those groups. This may allow us to better understand brain mechanisms that protect participants who share familiar risk but are unaffected.

## Acknowledgements

The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement n° 602450 (IMAGEMEND). In addition, the study was supported by grants from the Netherlands Organization for Scientific Research (NWO), i.e. a Vici grant to Franke (grant 016-130-669), Brain & Cognition grants 433-09-242 and 056-13-015 to Buitelaar, and the Gravitation Programme Language in Interaction (grant 024.001.006). The research of Franke and Buitelaar also receives funding from the FP7 Programme under grant agreements n° 602805 (AGGRESSOTYPE), n° 603016 (MATRICS) and n° 278948 (TACTICS), from the European Community's Horizon 2020 Programme (H2020/2014 – 2020) under grant agreements n° 643051 (MiND) and n° 642996 (BRAINVIEW), and from a grant for the ENIGMA Center for world-wide Medicine Imaging and Genomics from the National Institute of Health's BD2K Initiative (grant U54 EB020403). Andre Marquand gratefully acknowledges support from the Language in Interaction consortium, funded by the NWO under the Gravitation Programme.

**Financial disclosures**

Barbara Franke has received a speaker fee from Merz. Jan K Buitelaar has been in the past 3 years a consultant to / member of advisory board of / and/or speaker for Janssen Cilag BV, Eli Lilly, Shire, Lundbeck, Roche and Servier. He is not an employee of any of these companies, and not a stock shareholder of any of these companies. He has no other financial or material support, including expert testimony, patents, and royalties. Pieter Hoekstra has been member of the advisory board of Shire. The other authors do not report conflicts of interest. None of these, or any of the funding agencies have had any influence on the content of this manuscript.

## References

- Barkley, R.A., 1999. Response Inhibition in Attention-Deficit Hyperactivity Disorder. *Ment. Retard. Dev. Disabil. Res. Rev.* 5, 177–184. doi:10.1002/(SICI)1098-2779(1999)5:3<177::AID-MRDD3>3.0.CO;2-G
- Beckmann, C.F., Smith, S.M., 2004. Probabilistic independent component analysis for functional magnetic resonance imaging. *IEEE Trans. Med. Imaging* 23, 137–152.
- Biederman, J., Faraone, S.V., Keenan, K., Knee, D., Tsuang, M.T., 1990. Family-Genetic and Psychosocial Risk Factors in DSM-III Attention Deficit Disorder. *J. Am. Acad. Child Adolesc. Psychiatry* 29, 526–533. doi:10.1097/00004583-199007000-00004
- Birmaher, B., Ehmann, M., Axelson, D. a, Goldstein, B.I., Ph, D., Monk, K., Kalas, C., Kupfer, D., Gill, M.K., Leibenluft, E., Bridge, J., Guyer, A., Egger, H.L., Brent, D. a, 2010. Schedule for Affective Disorders and Schizophrenia for School- Age Children (K-SADS-PL) for the Assessment of Preschool Children- A Preliminary Psychometric Study. *Psychiatry (Abingdon)* 43, 680–686. doi:10.1016/j.jpsychires.2008.10.003.Schedule
- Castellanos, F.X., Sonuga-Barke, E.J.S., Milham, M.P., Tannock, R., 2006. Characterizing cognition in ADHD: beyond executive dysfunction. *Trends Cogn. Sci.* 10, 117–123. doi:10.1016/j.tics.2006.01.011
- Conners, K.C., Sitarenios, G., Parker, J.D.A., Epstein, J.N., 1998. The revised Conners' Parent Rating Scale (CPRS-R): Factor structure, reliability, and criterion validity. *J. Abnorm. Child Psychol.* 26, 257–268. doi:10.1023/A:1022602400621
- Durston, S., Mulder, M., Casey, B.J., Ziermans, T., van Engeland, H., 2006. Activation in ventral prefrontal cortex is sensitive to genetic vulnerability for attention-deficit hyperactivity disorder. *Biol. Psychiatry* 60, 1062–1070. doi:10.1016/j.biopsych.2005.12.020



- Efron, B., 2004. Large-scale simultaneous hypothesis testing: The choice of a null hypothesis. *J. Am. Stat. Assoc.* 99, 96–104. doi:10.1198/016214504000000089
- Efron, B., Tibshirani, R., Storey, J.D., Tusher, V., 2001. Empirical Bayes Analysis of a Microarray Experiment. *J. Am. Stat. Assoc.* 96, 1151–1160. doi:10.1198/016214501753382129
- Fair, D.A., Bathula, D., Nikolas, M.A., Nigg, J.T., 2012. Distinct neuropsychological subgroups in typically developing youth inform heterogeneity in children with ADHD. *Proc. Natl. Acad. Sci. U. S. A.* 109, 6769–74. doi:10.1073/pnas.1115365109
- Fan, Y., Gur, R.E., Gur, R.C., Wu, X., Shen, D., Calkins, M.E., Davatzikos, C., 2008. Unaffected Family Members and Schizophrenia Patients Share Brain Structure Patterns: A High-Dimensional Pattern Classification Study. *Biol. Psychiatry* 63, 118–124. doi:10.1016/j.biopsych.2007.03.015
- Faraone, S.V., Asherson, P., Banaschewski, T., Biederman, J., Buitelaar, J.K., Ramos-Quiroga, J.A., Rohde, L.A., Sonuga-Barke, E.J.S., Tannock, R., Franke, B., 2015. Attention-deficit/hyperactivity disorder. *Nat. Rev. Dis. Prim.* 1, 15020. doi:10.1038/nrdp.2015.20
- Faraone, S.V., Biederman, J., 2005. What is the prevalence of adult ADHD? Results of a population screen of 966 adults. *J. Atten. Disord.* 9, 384–391. doi:10.1177/1087054705281478
- Greven, C.U., Bralten, J., Mennes, M., O'Dwyer, L., van Hulzen, K.J.E., Rommelse, N., Schweren, L.J.S., Hoekstra, P.J., Hartman, C.A., Heslenfeld, D., Oosterlaan, J., Faraone, S.V., Franke, B., Zwiers, M.P., Arias-Vasquez, A., Buitelaar, J.K., 2015. Developmentally stable whole-brain volume reductions and developmentally sensitive caudate and putamen volume alterations in those with attention-deficit/hyperactivity disorder and their unaffected siblings. *JAMA psychiatry* 72,

490–9. doi:10.1001/jamapsychiatry.2014.3162

- Hart, H., Chantiluke, K., Cubillo, A.I., Smith, A.B., Simmons, A., Brammer, M.J., Marquand, A.F., Rubia, K., 2014a. Pattern classification of response inhibition in ADHD: Toward the development of neurobiological markers for ADHD. *Hum. Brain Mapp.* 35, 3083–3094. doi:10.1002/hbm.22386
- Hart, H., Marquand, A.F., Smith, A., Cubillo, A., Simmons, A., Brammer, M., Rubia, K., 2014b. Predictive neurofunctional markers of attention-deficit/hyperactivity disorder based on pattern classification of temporal processing. *J. Am. Acad. Child Adolesc. Psychiatry* 53, 569–578. doi:10.1016/j.jaac.2013.12.024
- Hastie, T., Tibshirani, R., Friedman, J., 2009. *The Elements of Statistical Learning*, 2nd ed. Springer. doi:10.1007/b94608
- Haufe, S., Meinecke, F., Görgen, K., Dähne, S., Haynes, J.-D., Blankertz, B., Bießmann, F., 2014. On the interpretation of weight vectors of linear models in multivariate neuroimaging. *Neuroimage* 87, 96–110. doi:10.1016/j.neuroimage.2013.10.067
- Ho, D.E., Imai, K., King, G., Stuart, E.A., 2011. MatchIt : Nonparametric Preprocessing for Parametric Causal Inference. *J. Stat. Softw.* 42, 1–28.
- Holm, S., 2010. A simple sequentially rejective multiple test procedure. *Statistics (Ber)*. 6, 65–70.
- Igual, L., Soliva, J.C., Escalera, S., Gimeno, R., Vilarroya, O., Radeva, P., 2012. Automatic brain caudate nuclei segmentation and classification in diagnostic of Attention-Deficit/Hyperactivity Disorder. *Comput. Med. Imaging Graph.* 36, 591–600. doi:10.1016/j.compmedimag.2012.08.002
- Janssen, T.W.P., Heslenfeld, D.J., Mourik, R.V., Logan, G.D., Oosterlaan, J., 2015. Neural Correlates of Response Inhibition in Children with Attention-Deficit/Hyperactivity Disorder: A Controlled Version of the Stop-Signal Task. *Psychiatry Res.*

- Neuroimaging 233, 1–7. doi:10.1016/j.psychresns.2015.07.007
- Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. Fsl. Neuroimage 62, 782–790. doi:10.1016/j.neuroimage.2011.09.015
- Johnston, B.A., Mwangi, B., Matthews, K., Coghill, D., Konrad, K., Steele, J.D., 2014. Brainstem abnormalities in attention deficit hyperactivity disorder support high accuracy individual diagnostic classification. Hum. Brain Mapp. 35, 5179–5189. doi:10.1002/hbm.22542
- Kottas, M., Kuss, O., Zapf, A., 2014. A modified Wald interval for the area under the ROC curve (AUC) in diagnostic case-control studies. BMC Med. Res. Methodol. 14, 26. doi:10.1186/1471-2288-14-26
- Lim, L., Marquand, A., Cubillo, A. a, Smith, A.B., Chantiluke, K., Simmons, A., Mehta, M., Rubia, K., 2013. Disorder-specific predictive classification of adolescents with attention deficit hyperactivity disorder (ADHD) relative to autism using structural magnetic resonance imaging. PLoS One 8, e63660. doi:10.1371/journal.pone.0063660
- Lipszyc, J., Schachar, R., 2010. Inhibitory control and psychopathology: a meta-analysis of studies using the stop signal task. J. Int. Neuropsychol. Soc. 16, 1064–1076. doi:10.1017/S1355617710000895
- Marquand, A., Howard, M., Brammer, M., Chu, C., Coen, S., Mourão-Miranda, J., 2010. Quantitative prediction of subjective pain intensity from whole-brain fMRI data using Gaussian processes. Neuroimage 49, 2178–89. doi:10.1016/j.neuroimage.2009.10.072
- Marquand, A.F., Rezek, I., Buitelaar, J., Beckmann, C.F., 2016a. Understanding heterogeneity in clinical cohorts using normative models: beyond case control studies. Biol. Psychiatry. doi:10.1016/j.biopsych.2015.12.023

- Marquand, A.F., Wolfers, T., Mennes, M., Buitelaar, J., Beckmann, C.F., 2016b. Beyond lumping and splitting: a review of computational approaches for stratifying psychiatric disorders. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging*. doi:10.1016/j.bpsc.2016.04.002
- Mostert, J.C., Hoogman, M., Onnink, A.M.H., van Rooij, D., von Rhein, D., van Hulzen, K.J.E., Dammers, J., Kan, C.C., Buitelaar, J.K., Norris, D.G., Franke, B., 2015a. Similar Subgroups Based on Cognitive Performance Parse Heterogeneity in Adults With ADHD and Healthy Controls. *J. Atten. Disord.* 1–12. doi:10.1177/1087054715602332
- Mostert, J.C., Onnink, A.M.H., Klein, M., Dammers, J., Harneit, A., Schulten, T., van Hulzen, K.J.E., Kan, C.C., Slaats-Willemse, D., Buitelaar, J.K., Franke, B., Hoogman, M., 2015b. Cognitive heterogeneity in adult attention deficit/hyperactivity disorder: A systematic analysis of neuropsychological measurements. *Eur. Neuropsychopharmacol.* 25, 2062–2074. doi:10.1016/j.euroneuro.2015.08.010
- Peng, X., Lin, P., Zhang, T., Wang, J., 2013. Extreme learning machine-based classification of ADHD using brain structural MRI data. *PLoS One* 8. doi:10.1371/journal.pone.0079476
- Polanczyk, G., de Lima, M.S., Horta, B.L., Biederman, J., Rohde, L.A., 2007. The worldwide prevalence of ADHD: a systematic review and metaregression analysis. *Am. J. Psychiatry* 164, 942–8. doi:10.1176/appi.ajp.164.6.942
- Pruim, R.H.R., Mennes, M., Buitelaar, J.K., Beckmann, C.F., 2015. Evaluation of ICA-AROMA and alternative strategies for motion artifact removal in resting state fMRI. *Neuroimage* 112, 278–287. doi:10.1016/j.neuroimage.2015.02.063
- Pruim, R.H.R., Mennes, M., van Rooij, D., Llera Arenas, A., Buitelaar, J.K., Beckmann, C.F., 2015. ICA-AROMA: A robust ICA-based strategy for removing motion artifact from

- fMRI data. *Neuroimage* 112, 267–277. doi:10.1016/j.neuroimage.2015.02.064
- Rasmussen, C.E., Williams, C.K.I., 2006. Model Selection and Adaptation of Hyperparameters. *Gaussian Process. Mach. Learn. (Adaptive Comput. Mach. Learn. Ser.* 105–128.
- Sabuncu, M.R., Konukoglu, E., 2014. Clinical Prediction from Structural Brain MRI Scans: A Large-Scale Empirical Study. *Neuroinformatics* 13, 31–46. doi:10.1007/s12021-014-9238-1
- Schrouff, J., Rosa, M.J., Rondina, J.M., Marquand, a F., Chu, C., Ashburner, J., Phillips, C., Richiardi, J., Mourão-Miranda, J., 2013. PRoNTTo: pattern recognition for neuroimaging toolbox. *Neuroinformatics* 11, 319–37. doi:10.1007/s12021-013-9178-1
- Segovia, F., Holt, R., Spencer, M., Górriz, J.M., Ramírez, J., Puntonet, C.G., Phillips, C., Chura, L., Baron-Cohen, S., Suckling, J., 2014. Identifying endophenotypes of autism: a multivariate approach. *Front. Comput. Neurosci.* 8, 1–8. doi:10.3389/fncom.2014.00060
- Simon, V., Czobor, P., Balint, S., Meszaros, A., Bitter, I., 2009. Prevalence and correlates of adult Attention-Deficit Hyperactivity Disorder: meta-analysis. *Br. J. Psychiatry* 194, 204–211. doi:10.1192/bjp.bp.107.048827
- Slaats-Willemse, D., Swaab-Barneveld, H., de Sonnevile, L., van der Meulen, E., Buitelaar, J., 2003. Deficient response inhibition as a cognitive endophenotype of ADHD. *J. Am. Acad. Child Adolesc. Psychiatry* 42, 1242–1248. doi:10.1097/00004583-200310000-00016
- Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E.J., Johansen-Berg, H., Bannister, P.R., De Luca, M., Drobnjak, I., Flitney, D.E., Niaz, R.K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J.M., Matthews, P.M., 2004. Advances

- in functional and structural MR image analysis and implementation as FSL. *Neuroimage* 23, 208–219. doi:10.1016/j.neuroimage.2004.07.051
- The ADHD Consortium, 2012. The ADHD-200 Consortium: A Model to Advance the Translational Potential of Neuroimaging in Clinical Neuroscience. *Front. Syst. Neurosci.* 6. doi:10.3389/fnsys.2012.00062
- Van Essen, D.C., Drury, H. a., Dickson, J., Harwell, J., Hanlon, D., Anderson, C.H., 2001. An Integrated Software Suite for Surface-based Analyses of Cerebral Cortex. *J. Am. Med. Informatics Assoc.* 8, 443–459. doi:10.1136/jamia.2001.0080443
- van Hulst, B.M., de Zeeuw, P., Durston, S., 2014. Distinct neuropsychological profiles within ADHD: a latent class analysis of cognitive control, reward sensitivity and timing. *Psychol. Med.* 1–11. doi:10.1017/S0033291714001792
- van Meel, C.S., Heslenfeld, D.J., Oosterlaan, J., Sergeant, J.A., 2007. Adaptive control deficits in attention-deficit/hyperactivity disorder (ADHD): The role of error processing. *Psychiatry Res.* 151, 211–220. doi:10.1016/j.psychres.2006.05.011
- van Rooij, D., Hartman, C.A., Mennes, M., Oosterlaan, J., Franke, B., Rommelse, N., Heslenfeld, D., Faraone, S.V., Buitelaar, J.K., Hoekstra, P.J., 2015a. Altered neural connectivity during response inhibition in adolescents with attention-deficit/hyperactivity disorder and their unaffected siblings. *NeuroImage Clin.* 7, 325–335. doi:10.1016/j.nicl.2015.01.004
- van Rooij, D., Hoekstra, P.J., Mennes, M., von Rhein, D., Thissen, A.J. a. M., Heslenfeld, D., Zwiers, M.P., Faraone, S.V., Oosterlaan, J., Franke, B., Rommelse, N., Buitelaar, J.K., Hartman, C. a., 2015b. Distinguishing Adolescents With ADHD From Their Unaffected Siblings and Healthy Comparison Subjects by Neural Activation Patterns During Response Inhibition. *Am. J. Psychiatry appi.ajp.* 2014.1. doi:10.1176/appi.ajp.2014.13121635

- von Rhein, D., Mennes, M., van Ewijk, H., Groenman, A.P., Zwiers, M.P., Oosterlaan, J., Heslenfeld, D., Franke, B., Hoekstra, P.J., Faraone, S.V., Hartman, C., Buitelaar, J., 2014. The NeuroIMAGE study: a prospective phenotypic, cognitive, genetic and MRI study in children with attention-deficit/hyperactivity disorder. Design and descriptives. *Eur. Child Adolesc. Psychiatry*. doi:10.1007/s00787-014-0573-4
- Wang, X., Jiao, Y., Lu, Z., 2011. Discriminative analysis of resting-state brain functional connectivity patterns of Attention-Deficit Hyperactivity Disorder using Kernel Principal Component Analysis, in: Eighth International Conference on Fuzzy Systems and Knowledge Discovery (FSKD). Ieee, pp. 1938–1941. doi:10.1109/FSKD.2011.6019911
- Wang, X., Jiao, Y., Tang, T., Wang, H., Lu, Z., 2013. Altered regional homogeneity patterns in adults with attention-deficit hyperactivity disorder. *Eur. J. Radiol.* 82, 1552–7. doi:10.1016/j.ejrad.2013.04.009
- Wolfers, T., Buitelaar, J.K., Beckmann, C.F., Franke, B., Marquand, A.F., 2015. From estimating activation locality to predicting disorder: a review of pattern recognition for neuroimaging-based psychiatric diagnostics. *Neurosci. Biobehav. Rev.* 57, 328–349. doi:10.1016/j.neubiorev.2015.08.001
- Woolrich, M.W., Jbabdi, S., Patenaude, B., Chappell, M., Makni, S., Behrens, T., Beckmann, C., Jenkinson, M., Smith, S.M., 2009. Bayesian analysis of neuroimaging data in FSL. *Neuroimage* 45, S173–S186. doi:10.1016/j.neuroimage.2008.10.055
- Yu, Y., Shen, H., Zhang, H., Zeng, L.-L., Xue, Z., Hu, D., 2013. Functional connectivity-based signatures of schizophrenia revealed by multiclass pattern analysis of resting-state fMRI from schizophrenic patients and their healthy siblings. *Biomed. Eng. Online* 12. doi:10.1186/1475-925X-12-10
- Zhu, C.-Z., Zang, Y.-F., Cao, Q.-J., Yan, C.-G., He, Y., Jiang, T.-Z., Sui, M.-Q., Wang, Y.-F., 2008.

Fisher discriminative analysis of resting-state brain function for attention-deficit/hyperactivity disorder. *Neuroimage* 40, 110–20.

doi:10.1016/j.neuroimage.2007.11.029

ACCEPTED MANUSCRIPT



Table 1: Demographic and clinical characteristics of complete sample

	Participants with ADHD		Unaffected siblings		Healthy controls		Sig.
N	184		103		128		
Males	128		41		60		
Females	56		62		68		
	Mean	SD	Mean	SD	Mean	SD	
ADHD symptoms <sup>a</sup>	12.94	2.90	0.75	1.28	0.36	0.90	ADHD > Siblings = Controls
Age	17.24	3.27	17.12	4.06	16.36	3.24	ADHD = Siblings = Controls
Age range	8 <-> 25		7 <-> 27		9 <-> 23		
Estimated IQ <sup>b</sup>	95.13	16.84	102.20	15.79	106.03	14.17	ADHD < Siblings = Controls
IQ range	55 <-> 138		65 <-> 144		58 <-> 141		
	Mean	SD	Mean	SD	Mean	SD	
SSRT (ms)	270.3	61.91	252.52	49.32	258.83	52.65	ADHD > Siblings = Controls
ICV (ms)	0.211	0.052	0.18	0.047	0.17	0.041	ADHD > Siblings = Controls
Errors (n)	6.45	7.89	4.05	5.29	3.45	4.31	ADHD > Siblings = Controls
Current Medication	107		4		0		
Comorbid ODD <sup>c</sup>	34		24		1		
Comorbid CD <sup>c</sup>	4		7		0		
Comorbid RD <sup>c</sup>	27		12		0		

Note: ADHD = attention deficit/hyperactivity disorder; ODD = oppositional defiant disorder; CD = conduct disorder; RD = Reading disability; SSRT = Stop-signal reaction time; RTV = reaction time variability; Errors = number of errors on go-trials; Sig.= Nominal significant differences are listed in this column if this column is empty no significant differences could be reported.

<sup>a</sup> ADHD diagnosis was based on K-SADS structured psychiatric interviews and Conners' questionnaires (Conners et al., 1998).

<sup>b</sup> Estimated IQ was based on the block-design and vocabulary subtests of the Wechsler Intelligence Scale for Children (WISC) or Wechsler Adult Intelligence Scale (WAIS-III; Wechsler, 2012).

<sup>c</sup> ODD, CD, and RD diagnosis was based on K-SADS structured psychiatric interviews (Kaufman et al., 1997).

*Figure 1*

Depicted are the results for all predictions in the complete and matched sample, the x-axis corresponds to the area under the receiver-operating characteristic curve (AUC). An area of 0.5 indicates no discrimination; \*\* indicates that the prediction remains significant after Bonferroni-Holm correction; \* indicates that predictions are significant before multiple comparison correction.

*Figure 2*

Forward maps of significant predictions based on successful stop activation maps. **I)** Unaffected siblings versus healthy controls show a frontolateral pattern with positive coefficients favouring siblings. **II)** Unaffected siblings versus participants with ADHD show a widespread pattern with positive coefficients favouring ADHD predictions. **III)** Participants with ADHD versus healthy controls show a relatively clear frontolateral, interior-parietal and superior-temporal pattern with positive coefficients favouring ADHD predictions. The left images correspond to the left hemisphere, the right images to the right hemisphere.

*Supplementary Figure 1*

First column shows the forward maps of significant predictions based on successful stop activation maps. Second column shows Z-statistics thresholded at  $p_{\text{AHT}} > .5$  for the forward maps. Third column shows Z-statistics thresholded at  $p_{\text{FDR}} < .05$  the forward maps.

*Supplementary Figure 2*

Fit of the mixture models to the three forward maps depicted.

Figure 1





Figure 2



**Highlights**

- 1) Multivariate profile of risk shared between ADHD patients and unaffected siblings
- 2) Pattern of risk in frontolateral, superior temporal and inferior parietal regions
- 3) Benchmark estimate of classification performance in heterogeneous samples